

PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

NAME: _____

DATE: _____

Over the *last 2 weeks*, how often have you been bothered by any of the following problems?
(use "✓" to indicate your answer)

	Not at all 0	Several days 1	More than half the days 2	Nearly every day 3
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite—being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead, or of hurting yourself in some way	0	1	2	3

add columns: + +

(Healthcare professional: For interpretation of TOTAL, TOTAL: _____
please refer to accompanying scoring card).

10. If you checked off *any* problems, how *difficult* have these problems made it for you to do your work, take care of things at home, or get along with other people?

- Not difficult at all _____
- Somewhat difficult _____
- Very difficult _____
- Extremely difficult _____

PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

NAME: John Q. Sample

DATE: _____

Over the *last 2 weeks*, how often have you been bothered by any of the following problems?
(use "✓" to indicate your answer)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	✓	3
2. Feeling down, depressed, or hopeless	0	✓	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	✓	3
4. Feeling tired or having little energy	0	1	2	✓
5. Poor appetite or overeating	0	✓	2	3
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	0	1	✓	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	✓	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite—being so fidgety or restless that you have been moving around a lot more than usual	0	1	✓	3
9. Thoughts that you would be better off dead, or of hurting yourself in some way	✓	1	2	3

add columns: **2** + **10** + **3**

(Healthcare professional: For interpretation of TOTAL, TOTAL: **15**
please refer to accompanying scoring card).

<p>10. If you checked off <i>any</i> problems, how <i>difficult</i> have these problems made it for you to do your work, take care of things at home, or get along with other people?</p>	<p>Not difficult at all _____</p> <p>Somewhat difficult <u>✓</u> _____</p> <p>Very difficult _____</p> <p>Extremely difficult _____</p>
---	---

**INSTRUCTIONS
PHQ -9 SCORING TOOL**

Score	Action
≤ 4	The score suggests that patient may not need depression treatment
$\geq 5 - 14$	Physician uses clinical judgment about treatment, based on patient's duration of symptoms and functional impairment.
≥ 15	Warrants treatment for depression, using antidepressant psychotherapy and/or a combination of treatment

A functional health assessment is reflected in Question 3 on the PHQ-9, which asks the patient how emotional difficulties or problems impact work, things at home or relationships with other people. Patient responses can be one of four: not difficult at all, somewhat difficult, very difficult, or extremely difficult. A response of very difficult or extremely difficult suggests that the patient's functionality is impaired.